Major Land Resource Area 118A Arkansas Valley and Ridges, Eastern Part

Accessed: 05/10/2025

Ecological site keys

MLRA 118A ES Key

- I. Site receives additional moisture from the surrounding landscape (flood plains, terraces, and foot slopes), soils formed in alluvium.
 - A. Rare (1 to 5 times in 100 years) to frequent (greater than 50 times in 100 years) flooding occurs.
 - 1 Occurs on flood plains.
 - i. Soils are well to moderately well drained and characterized by an irregular decrease in organic carbon, stratified textures and/or a buried A horizon (fluventic soils). ... NX118A01Y008 Fluventic Flood Plain
 - ii. Soils are somewhat poorly to poorly drained and characterized by greater than 60 percent clay in the particle size control section and a layer of clay accumulation 20 to 30 cm from the surface. ... NX118A01Y004 Clayey Flood Plain
 - 2 Occurs on paleoterraces or high terraces and are subject to rare flooding for brief periods (2- less than 7 days).
 - i. Soils are well to moderately well drained and characterized by a layer of clay accumulation within 25 to 50 cm from the soil surface. ... NX118A01Y003 Rarely Flooded Terrace
 - B. Flooding does not occur.
 - 1 Soils are well to moderately well drained and characterized by a layer of clay accumulation within 25 to 50 cm from the soil surface. ... NX118A01Y007 Seasonally Wet Terraces and Footslopes
- II. Site does not receive additional moisture from the surrounding landscape (side slopes, plateaus, and ridges), soils formed in residuum and colluvium.
 - A. Soils are less than 50 cm to restrictive layer (shallow). ... NX118A01Y001 Shallow Upland
 - B. Soil greater than or equal to 50 cm to restrictive layer (moderately deep, deep, or very deep).
 - 1 Occurs on linear to convex backslope or side slopes of paleoterraces, plateaus, and ridges.
 - i. Greater than 35 percent clay (fine) in the particle size control section with a layer of clay accumulation within 25 cm of the soil surface. ... NX118A01Y005 Clayey Upland
 - ii. Less than 35 percent clay (fine-loamy or fine-silty) in particle size control section with a layer of clay accumulation within 10 to 35 cm from the soil surface. ... NX118A01Y002 Loamy Upland
 - 2 Occurs on gently sloping to concave uplands.
 - i. Greater than 35 percent clay (fine) in the particle size control section and moderately to strongly alkaline (pH 7.9 to 9.0) below 25 cm. ... NX118A01Y006 Alkali Flat